arXiv:1201.6068v2 [math.GT] 27 Jun 2012

COMPUTING THE RHO-INVARIANTS OF LINKS VIA THE SIGNATURE OF COLORED LINKS WITH APPLICATIONS TO THE LINEAR INDEPENDENCE OF TWIST KNOTS.

CHRISTOPHER WILLIAM DAVIS

Abstract. We use a link invariant defined by Cimasoni-Florens to compute certain ρ-invariants. This generalizes results of Cochran-Orr-Teichner and Friedl on the ρ-invariants of knots to the setting of links. As an application, we prove with only twelve possible exceptions that the twist knots of algebraic order two are linearly independent in the topological concordance group.

1. Introduction

A link is an isotopy class of smooth, ordered, oriented embeddings of a disjoint union of circles $S^1 \sqcup \cdots \sqcup S^1$ into the 3-sphere S^3. A knot can be thought of as a one component link. Two n-component links L and J are called **concordant** if there is a locally flat, ordered, oriented embedding of a disjoint union of n annuli $\sqcup(S^1 \times [0,1])$ into $S^3 \times [0,1]$ sending $\sqcup(S^1 \times \{0\})$ to L in $S^3 \times \{0\}$ and $\sqcup(S^1 \times \{1\})$ to J in $S^3 \times \{1\}$. If one restricts to knots then the set of concordance classes forms a group, $\mathcal{C}$, called the knot concordance group, under the operation of connected sum.

From a link L with zero pairwise linking numbers one can get a closed oriented 3-manifold, $M(L)$ by taking zero framed surgery. Given such a closed, oriented 3-manifold M and a unitary representation $\alpha : \pi_1(M) \to U(n)$ the unitary ρ-invariant $\rho(M, \alpha) \in \mathbb{Z}$ is defined [1]. Similarly, given a homomorphism to any group, $\phi : \pi_1(M) \to G$, the L^2-ρ-invariant, $\rho^{(2)}(M, \phi) \in \mathbb{R}$ is defined, see for example [14]. The ρ-invariants of $M(L)$ have been important tools in the study of concordance. For example, see [7], [6], [11], [17], [14] or [10].

These invariants are notoriously difficult to compute, although in the case of knots and representations to $U(1)$ Litherland [20] computes them in terms of the Tristram-Levine signature function. In the case knots and of homomorphisms to cyclic groups, Cochran-Orr-Teichner [8, Theorem 5.1] and Freidl [12, Corollary 4.3] show that L^2-ρ-invariants can be computed in terms of sums and integrals of the Tristram-Levine signature function.

Date: June 28, 2012.

2010 *Mathematics Subject Classification.* 57N65.

The main goal of this paper is the extension of these results computing one dimensional unitary ρ-invariants and abelian L^2-ρ-invariants of knots to tools to the setting of links. In Theorem 3.5, we find that for a link L with zero pairwise linking the ρ-invariants of $M(L)$ corresponding to representations to $U(1)$ are given by the evaluation of the signature function σ_L defined by Cimasoni-Florens in [4]. In Theorem 4.3 we express L^2-ρ-invariants corresponding to homomorphisms to abelian groups in terms of its integral. In [10], L^2-ρ-invariants of links are used to obstruct linear dependence of concordance classes of knots. In [10] an ad hoc method is used to compute these invariants and find an infinite family of twist knots (see Figure 1) which is linearly independent in the knot concordance group. We use Theorem 4.3 in order to prove that with only twelve possible exceptions the twist knots which are algebraically of order 2 (depicted in Figure 1) are linearly independent.

Theorem 5.5. *The set containing all of the twist knots T_n which are algebraically of order 2 is linearly independent with the following* 12 *possible exceptions:*

$$n = 1, 11, 16, 29, 36, 38, 51, 55, 61, 66, 83, 101.$$



FIGURE 1. The n-twist knot.

1.1. **Background: The Cimasoni-Florens signature function.** We now provide some of the background needed in this paper. We begin with a brief discussion of the signature function associated to colored links by Cimasoni and Florens in [4].

An n**-colored link** $L = L(1), \dots, L(n)$ is a link whose every component is decorated with an integer between 1 and n. The notation $L(k)$ refers to the sublink of L consisting of components colored with the integer k. Note that any n-component link $L_1, \dots, L_n$ has an n-coloring given by taking $L(n) = L_n$. Except when explicitly stated we will consider any link to be colored in this manner.

Let $\mathbb{T}^n$ denote the n-dimensional unit torus in $\mathbb{C}^n$,

$$\mathbb{T}^n := \{(\omega_1, \dots, \omega_n) \in \mathbb{C}^n : |\omega_k| = 1 \text{ for all } k\}.$$

The Cimasoni-Florens signature of an n-colored link L, σ_L is an integer valued function on

$$\mathbb{T}^n_* := \{(\omega_1, \dots, \omega_n) \in \mathbb{T}^n : \omega_k \neq 1 \text{ for all } k\}.$$

Its definition is in terms of the following generalization of Seifert surfaces to the context of colored links

Definition 1.1 (Subsecton 2.1 of [4]). A union of smoothly embedded transversely intersecting surfaces $S = S_1, \dots, S_n$ is a **C-complex** (or **Clasp-Complex**) for the n-colored link, $L = L(1) \dots L(n)$, if the following conditions hold:

1. For each i S_i is a Seifert surface for $L(i)$. That is, S_i is a compact oriented embedded surface with no closed components whose oriented boundary is $L(i)$.
2. For $i \neq j$, $S_i \cap S_j$ is a union of arcs each having one boundary point in $\partial S_i = L(i)$ and the other in $\partial S_j = L(j)$. These intersections are called **clasps**. See the leftmost part of Figure 2.
3. For distinct numbers i, j, k, $S_i \cap S_j \cap S_k$ is empty.



FIGURE 2. Left to Right: (1) A clasp in a C-Complex. (2) A loop γ passing through a clasp. (3) The curve γ^{+-} is given by pushing γ in the positive normal direction on one component on the C-complex and in the negative normal direction of the other.

It is worth noting that every colored link has a C-complex. In [4], Cimasoni and Florens define a matrix whose entries are Laurent polynomials in terms of a C-complex. We present an abbreviated version of their construction.

A simple closed curve on a C-complex F is called a **loop** if, whenever it intersects a clasp, it does so as in the second picture in Figure 2. Let $B = \{\gamma_1, \dots, \gamma_m\}$ be a basis for $H_1(F)$ given by loops.

For a loop $\gamma_j \in B$ and $\epsilon = \langle \epsilon_1 \dots \epsilon_n \rangle \in \{\pm 1\}^n$ define γ_j^ϵ by pushing the portion of γ_j carried by S_i off of S_i in the positive normal direction if $\epsilon_i = +1$ and in the negative normal direction otherwise (see the third picture in Figure 2). Let A^ϵ be the matrix whose (i,j)-entry is the linking number $\operatorname{lnk}(\gamma_i^\epsilon, \gamma_j)$. Define $H(\omega_1, \dots, \omega_n)$ by

$$H(\omega_1, \dots, \omega_n) = \sum_{\epsilon \in \{\pm 1\}^n} (1 - \omega_1^{-\epsilon_1}) \dots (1 - \omega_n^{-\epsilon_n}) A^\epsilon. \tag{1.1}$$

The matrix $H(\omega)$ is Hermitian for each $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{T}^n$. They define the following isotopy invariant.

Definition 1.2 (Subsecton 2.2 of [4]). **The Cimasoni-Florens signature function** of the n-colored link L, $\sigma_L : \mathbb{T}^n_* \to \mathbb{Z}$ is defined at $\omega \in \mathbb{T}^n_*$ to be the signature of $H(\omega)$, that is,

$$\sigma_L(\omega) := \sigma(H(\omega)) = \#\{\text{positive eigenvalues}\} - \#\{\text{negative eigenvalues}\}.$$

In [4], Cimasoni and Florens prove that σ_L is independent of the C-Complex F and the basis B, that is, σ_L is a color preserving isotopy invariant of L.

1.2. **Signatures and ρ-invariants.** We begin this subsection with a hasty review of signature invariants of 4-manifolds. Let W be a compact oriented 4-manifold. The classical signature, $\sigma(W)$, is defined in terms of a symmetric bilinear form $Q_W : H_2(W;\mathbb{Q}) \times H_2(W;\mathbb{Q}) \to \mathbb{Z}$ called the intersection form. By diagonalizing Q_W, the second homology of W is seen to decompose as a direct sum $H_2^+(W;\mathbb{Q}) \oplus H_2^-(W;\mathbb{Q}) \oplus H_2^0(W;\mathbb{Q})$ such that Q_W is positive definite on $H_2^+(W;\mathbb{Q})$, negative definite on $H_2^+(W;\mathbb{Q})$, and zero on $H_2^0(W;\mathbb{Q})$. The signature of W is given by

$$\sigma(W) = \sigma(Q_W) = \operatorname{rank}(H_2^+(W;\mathbb{Q})) - \operatorname{rank}(H_2^-(W;\mathbb{Q})).$$

The twisted and L^2-signatures of W are defined in terms of the twisted and L^2-homology groups, which we now recall. Let $\alpha : \pi_1(W) \to U(n)$ be a unitary representation of the fundamental group of W and $\phi : \pi_1(W) \to \Gamma$ be a group homomorphism. From this the vector space $\mathbb{C}^n$ and the Hilbert space $l^2(\Gamma)$ are given the structure of $\mathbb{Z}[\pi_1(W)]$ modules. We encode the module structure of $\mathbb{C}^n$ with the notation $\mathbb{C}^n_\alpha$. The cellular chain groups of the universal cover of W, $C_*(\widetilde{W})$ also have the structure of a $\mathbb{Z}[\pi_1(W)]$-module and the boundary maps are module homomorphisms.

Consider the tensored chain complexes:

$$\left(C_*(W;\mathbb{C}^n_\alpha), \partial^\alpha_*\right) = \left[\xrightarrow{\partial_{k+1}\otimes 1} C_k(\widetilde{W}) \otimes \mathbb{C}^n_\alpha \xrightarrow{\partial_k \otimes 1} C_{k-1}(\widetilde{W}) \otimes \mathbb{C}^n_\alpha\right],$$

$$\left(C^{(2)}_*(W;l^2(\Gamma)), \partial^{(2)}_*\right) = \left[\xrightarrow{\partial_{k+1}\otimes 1} C_k(\widetilde{W}) \otimes l^2(\Gamma) \xrightarrow{\partial_k \otimes 1} C_{k-1}(\widetilde{W}) \otimes l^2(\Gamma)\right].$$

The homology groups of W are given by the homology of $(C_*(W;\mathbb{C}^n_\alpha), \partial_*)$,

$$H_k(W;\mathbb{C}^n_\alpha) = \frac{\ker(\partial^\alpha_k)}{\operatorname{im}(\partial^\alpha_{k+1})}$$

while the L^2-homology groups are given by

$$H^{(2)}_k(W;l^2(\Gamma)) = \frac{\ker(\partial^{(2)}_k)}{\operatorname{clos}(\operatorname{im}(\partial^{(2)}_{k+1}))}$$

where $\operatorname{clos}(\operatorname{im}(\partial^{(2)}_{k+1}))$ denotes the closure of the subspace $\operatorname{im}(\partial^{(2)}_{k+1})$ of the Hilbert space, $C^{(2)}_k(W;l^2(\Gamma))$.

There are Hermitian bilinear forms on the twisted second homology $H_2(W;\mathbb{C}^n_\alpha)$ and the L^2-second homology $H^{(2)}_2(W;l^2[\Lambda])$ called twisted and L^2-intersection forms. They are denoted Q^α_W and $Q^{(2)}_W$. By diagonalizing Q^α_W and applying the spectral theorem to $Q_W^{(2)}$, these homology groups are seen to decompose into subspaces on which the associated intersection forms are positive definite, negative definite and zero:

$$\begin{array}{rcl} H_2(W,\mathbb{C}_\alpha) & \cong & H_2^+(W,\mathbb{C}_\alpha)\oplus H_2^-(W,\mathbb{C}_\alpha)\oplus H_2^0(W,\mathbb{C}_\alpha) \\ H_2^{(2)}(W;l^2(\Gamma)) & \cong & H_2^{(2)+}(W;l^2(\Gamma))\oplus H_2^{(2)-}(W;l^2(\Gamma))\oplus H_2^{(2)0}(W;l^2(\Gamma)) \end{array}$$

The twisted signature invariant is given by

$$\sigma(W,\alpha)=\dim(H_2^+(W,\mathbb{C}_\alpha))-\dim(H_2^-(W,\mathbb{C}_\alpha)),$$

while the L^2-signature invariant is given by

$$\sigma^{(2)}(W,\phi)=\dim_{\mathcal{N}\Lambda}(H_2^{(2)+}(W,l^2[\Lambda]))$$

where dim denotes the classical dimension of $\mathbb{C}$-vector spaces and $\dim_{\mathcal{N}\Lambda}$ denotes the von Neumann dimension of projective $\mathcal{N}\Lambda$-modules. For more information on the von Neumann dimension and the L^2-signature invariant the reader is referred to [22] and [23].

In [1], Atiyah, Patodi and Singer associate to a 3-manifold M and a unitary representation $\alpha:\pi_1(M)\to U(n)$ an integer, $\rho(M,\alpha)$. This invariant has the property that if W is a 4-manifold with a unitary representation $\alpha:\pi_1(W)\to U(n)$ and boundary components, $M_1\ldots M_k$ then

$$\sum_{i=1}^{k}\rho(M_i,\alpha_i)=\sigma(W,\alpha)-n\sigma(W)$$

where α_i is given by the composition $\pi_1(M_i)\to\pi_1(W)\overset{\alpha}{\to}U(n)$.

For a closed oriented 3-manifold M with coefficient system $\phi:\pi_1(M)\to\Gamma$ the von Neumann $\rho^{(2)}$-invariant is given in terms of a coefficient system on a bounded 4-manifold, for example see [14, Section 3]. More precisely, for 3-manifolds $M_1,\ldots,M_k$ with coefficient systems $\phi_i:\pi_1(M_i)\to\Gamma_i$, if W is a compact oriented connected 4-manifold with $\partial W=\bigsqcup_{i=1}^{n}M_i$ and $\psi:\pi_1(W)\to\Lambda$ is a coefficient system on W such that such that for each i there is a monomorphism $\theta_i:\Gamma_i\hookrightarrow\Lambda$ making the following diagram commute:

(1.2)
$$\begin{tikzcd} \pi_1(M_i) \arrow[r, "\phi_i"] \arrow[d, "i_*"] & \Gamma_i \arrow[d, hook, "\theta_i"] \\ \pi_1(W) \arrow[r, "\psi"] & \Lambda \end{tikzcd}$$

Then $\sum_{i=1}^{k}\rho^{(2)}(M_i,\phi_i)=\sigma^{(2)}(W,\psi)-\sigma(W)$.

1.3. **Statement of results.** Before we can state the results of this paper, we need some additional notation. For an integrer n, $\mathbb{T}^n_{\mathbb{Q}}$ refers to the set of n-tuples of roots of unity:

$$\mathbb{T}^n_{\mathbb{Q}} = \{(z_1, \dots, z_n) \in \mathbb{T}^n : z_i \text{ is a root of unity for } i = 1, \dots n\}.$$

Given any finitely generated abelian group $A = \langle g_1, \dots, g_n | r_1 \dots r_m \rangle$ where $r_i = \sum_{j=1}^{n} a_{i,j} g_j$ for integers $a_{i,j}$, we also define

$$\mathbb{T}_A = \left\{ (z_1, \dots, z_n) : \prod_{j=1}^{n} z_j^{a_{i,j}} = 1 \text{ for all } i \right\}. \tag{1.3}$$

The space $\mathbb{T}_A$ is isomorphic to the space of one dimensional representations of A. The isomorphism is given by sending $z = (z_1, \dots, z_n) \in \mathbb{T}^n$ to the representation $A \to U(1)$ sending g_i to the one-by-one matrix, $[z_i]$.

In this paper we prove the following theorems.

Theorem 3.5. *For a link L with zero pairwise linking there is a piecewise constant function $\widehat{\sigma_L} : \mathbb{T}^n \to \mathbb{Z}$ which agrees with σ_L on $\mathbb{T}^n_*$. For $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{T}^n_{\mathbb{Q}}$ and the representation $\alpha_\omega : \pi_1(M(L)) \to U(1)$ given by sending the meridian, μ_i, to ω_i, $\rho(M(L), \alpha_\omega) = \widehat{\sigma}_L(\omega)$.*

Theorem 4.3. *Let $A = \langle g_1 \dots g_n | r_1 \dots r_m \rangle$ be an abelian group. Let L be an n-component link with zero pairwise linking numbers. Let $\phi : H_1(M(L)) \to A$ be given by sending μ_i to g_i, then*

$$\rho^{(2)}(M(L), \phi) = \frac{1}{\lambda(\mathbb{T}_A)} \int_{\mathbb{T}_A} \widehat{\sigma}_L(\omega) d\lambda(\omega)$$

where λ is Lebesgue measure on $\mathbb{T}_A$.

Remark 1.3. We provide this extension $\widehat{\sigma}$ of the Cimasoni-Florens signature in Definition 3.4. For now we will merely comment that this function will remain quite computable.

One connection between Theorem 4.3 and knot concordance comes in the results of [10]. We use Theorem 4.3 to approximate the obstruction to linear dependence in $\mathcal{C}$ which appears in [10] for the twist knots that are algebraically of order 2. In doing so we prove Theorem 5.5 showing that with only finitely many exceptions those twist knots which are of order 2 in the algebraic concordance group are linearly independent in the topological concordance group.

In previous results, Livingston and Naik in [21] showed that infinitely many of those twist knots that are algebraically of order 4 are not of finite order in $\mathcal{C}$. Tamulis [27] found an infinite linearly independent set of algebraic order 2 twist knots. Cha [16] showed that no nontrivial linear combination of the twist knots (with the exception of the unknot, stevedore knot and figure eight knot) is ribbon. Results of Lisca, [19], show that the twist knots (with the same exceptions) are linearly independent in the *smooth* concordance group.

2. The 4-manifold used to compute abelian ρ-invariants of links

Consider any link, L, with zero pairwise linking numbers. In this section we build a 4-manifold W with $M(L)$ as a boundary component such that the inclusion induced map $H_1(M(L)) \to H_1(W)$ is an isomorphism. Since all one dimensional unitary representations of $\pi_1(M(L))$ and all homomorphisms from $\pi_1(M(L))$ to an abelian group factor through $H_1(M(L))$ they extend over $\pi_1(W)$. Thus, we will be able to compute one dimensional unitary and abelian ρ-invariants of $M(L)$ in terms of the signature defects of W.

Construction 2.1. For an n-component link $L = L_1 \dots L_n$ with zero pairwise linking, let $F = F_1 \dots F_n$ be a collection of n embedded disjoint surfaces in B^4 such that $\partial F_i = L_i$. Such a collection exists because L has zero pairwise linking. Let $E(F)$ denote the exterior of F in B^4. The exterior of L, $E(L)$, is a submanifold of $\partial E(F)$. To a neighborhood of each component of the boundary of $E(L)$ (a torus in S^3 enclosing a component of L) attach to $E(F)$ a copy of $S^1 \times B^2 \times I$ so that the nullhomologous longitudes of the components of L bound a disks. Call the resulting 4-manifold W.

The boundary of W consists of a copy of $M(L)$ together with the orientation reverse of $\widehat{F} \times S^1$. The n-component closed surface $\widehat{F} := \widehat{F_1} \sqcup \dots \sqcup \widehat{F_n}$ is given by capping the only boundary component of each F_i with a disk.

The following lemma will be used to compute the ρ-invariants of $M(L)$:

Lemma 2.2. *Let W be the 4-manifold given in Construction 2.1. Then*

1. *the inclusion induced map $H_1(M(L)) \to H_1(W)$ is an isomorphism;*
2. $\sigma(W) = 0$.

Before proving Lemma 2.2 we explore its uses. As a consequence, we have the following commutative diagram for any one dimensional unitary representation $\alpha : \pi_1(M(L)) \to U(1)$. Since $U(1)$ is abelian we may as well regard α as a representation of $H_1(M(L))$.

$$\begin{tikzcd}
\pi_1(M(L)) \arrow[r] \arrow[d] & H_1(M(L)) \arrow[r, "\alpha"] \arrow[d, "\cong"'] & U(1) \\
\pi_1(W) \arrow[r] & H_1(W) \arrow[ur, "\alpha"] &
\end{tikzcd}$$

It follows that

$$\rho(M(L),\alpha) - \sum_{i=1}^{n} \rho(\widehat{F_i} \times S^1, \alpha) = \sigma(W,\alpha) - \sigma(W) = \sigma(W,\alpha). \tag{2.1}$$

Thus, in order to prove Theorem 3.5 , it will suffice to analyze the twisted signatures of W and the one dimensional unitary ρ-invariants of product

3-manifolds. This is done is Section 3. Similarly, any homomorphism to an abelian group $\phi : \pi_1(M(L)) \to A$ extends over W and

$$\rho^{(2)}(M(L),\phi) - \sum_{i=1}^{n} \rho^{(2)}(\widehat{F}_i \times S^1,\phi) = \sigma^{(2)}(W,\phi). \tag{2.2}$$

In Section 4 we compute $\sigma^{(2)}(W,\phi)$ and $\rho^{(2)}(\widehat{F}_i \times S^1,\phi)$. In doing so we prove Theorem 4.3.

Proof of Lemma 2.2. In [4, Section 6] an argument using the long exact sequence of the pair $(B^4, E(F))$ and duality shows that the inclusion induced map $H_1(E(L)) \to H_1(E(F))$ is an isomorphism. Since the longitudes are nullhomologous, it follows that the inclusion induced maps $H_1(E(L)) \to H_1(M(L))$ and $H_1(E(F)) \to H_1(W)$ are isomorphisms. It follows that the bottom row of the following commutative diagram is an isomorphism

$$\begin{tikzcd}
H_1(E(L)) \arrow[r, "\cong"] \arrow[d, "\cong"'] & H_1(E(F)) \arrow[d, "\cong"'] \\
H_1(M(L)) \arrow[r] & H_1(W)
\end{tikzcd}$$

In order to see (2), first observe that if A and B are classes in $H_2(E(F))$ whose intersection pairing $Q_{E(F)}(A,B)$ is nonzero and i_* denotes the inclusion induced map $H_2(E(F)) \to H_2(B^4)$, then

$$Q_B(i_*(A), i_*(B)) = Q_{E(F)}(A,B) \neq 0.$$

Which is a contradiction, since $H_2(B^4) = 0$. The intersection form $Q_{E(F)}$ must be zero. Since the longitudes of L are nullhomologous $H_2(W) = H_2(E(F)) \oplus \mathbb{Z}^n$. The $\mathbb{Z}^n$-factor has basis given by the embedded surfaces $\widehat{F}_i$. These embedded surfaces sit in the boundary of W and so have zero intersection numbers. Thus, the intersection matrix for W is a zero matrix, $H_2^+(W) = H_2^-(W) = 0$, and $\sigma(W) = 0$. □

3. Computing unitary ρ-invariants

In this section we prove Theorem 3.5. In light of equation (2.1) it remains only to analyze the one dimensional unitary signatures of W and the one dimensional unitary ρ-invariants of the product 3-manifold $F \times S^1$

First we recall a result of Cimasoni-Florens [4]. As stated and proven in [4], it requires neither zero linking numbers nor that all components have distinct colors.

Theorem ([4], Theorem 6.1). *Let $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{T}^n_{\mathbb{Q}} \cap \mathbb{T}^m_*$ be an n-tuple of roots of unity none of which are equal to 1. Let $L = L_1, \dots, L_n$ be an n-component link with zero pairwise linking. Regard L as an n-colored link by coloring each component differently. For a union of n disjoint, compact,* *connected, oriented, properly embedded surfaces* $F = F_1 \cup \cdots \cup F_n \subseteq B^4$ *with* $\partial F_i = L_i$ *and the representation* $\alpha_\omega : \pi_1(E(F)) \to U(1)$ *sending* μ_i*, the meridian of* L_i*, to* $[\omega_i]$*,* $\sigma(E(F), \alpha_\omega) = \sigma_L(\omega)$*.*

We prove the following propositions:

Proposition 3.1. *For the 4-manifold* W *of Construction 2.1 and* $\omega \in \mathbb{T}^n_*$*, the inclusion induced map* $H_2(E(F); \mathbb{C}_{\alpha_\omega}) \to H_2(W; \mathbb{C}_{\alpha_\omega})$ *is an isomorphism so that* $\sigma(W, \alpha_\omega) = \sigma(E(F), \alpha_\omega)$*.*

Proposition 3.2. *For a closed oriented connected surface* F*, and any representation* $\alpha : \pi_1(F \times S^1) \to U(1)$ *with image in the roots of unity,* $\rho(F \times S^1, \alpha) = 0$

Putting this all together, for any $\omega \in \mathbb{T}^n_{\mathbb{Q}} \cap \mathbb{T}^n_*$

(3.1)
$$\begin{array}{rcll} \rho(M(L), \alpha_\omega) & = & \sigma(W, \alpha_\omega) - \sum_i \rho(\widehat{F_i} \times S^1, \alpha_\omega) & \text{by equation (2.1)} \\ & = & \sigma(W, \alpha_\omega) & \text{by Proposition 3.2} \\ & = & \sigma(E(F), \alpha_\omega) & \text{by Proposition 3.1} \\ & = & \sigma_L(\omega) & \text{by [4, Theorem 6.1]} \end{array}$$

This proves Theorem 3.5 in the case that no ω_i is equal to 1. The main difficulty in completing the proof is that the Cimasoni-Florens signature is only defined when $\omega_i \neq 1$ for all i. As the following theorem indicates, even in this setting the Cimasoni-Florens signature (of a different colored link) computes these unitary ρ-invariants.

Proposition 3.3. *Let* $L = L_1 \dots L_n$ *be an* n*-component link with zero pairwise linking. Let* $\omega \in \mathbb{T}^n_{\mathbb{Q}}$ *be an* n*-tuple of roots of unity. Let* $A = \{k : \omega_k = 1\}$ *and* $m = |A|$ *be the cardinality of* A*. Let* J *be the* n*-colored,* $(n+m)$*-component link given by replacing* L_k *by two zero framed pushoffs of* L_k *with opposite orientations and the same color for each* $k \in A$*. Let*

$$\omega^J = (\omega^J_1, \dots, \omega^J_n) \in \mathbb{T}^n_{\mathbb{Q}} \cap \mathbb{T}^n_*$$

be any n*-tuple of roots of unity not equal to 1 with* $\omega^J_k = \omega_k$ *if* $k \notin A$*.*

Then $\rho(M(L), \alpha_\omega) = \sigma_J(\omega^J)$

Proof. Let $L^0 = L_1, \dots, L_n, U_1, \dots, U_m$ be the $(n+m)$-color, $(n+m)$-component link given by the split union of L with an m-component unlink.

Let ω^0 be an $(n+m)$-tuple of roots of unity given by $(\omega_1, \dots, \omega_n, z_1, \dots, z_m)$ where $(z_1, \dots, z_m) \in \mathbb{T}^m_{\mathbb{Q}} \cap \mathbb{T}^m_*$ is an m-tuple of roots of unity other than 1. Since $M(L^0) = M(L) \#(\#_m S^2 \times S^1)$, ρ-invariants are additive over connected sum of 3-manifolds and the one dimensional unitary ρ-invariants of $S^2 \times S^1$ vanish (for example, by Proposition 3.2), it follows that

$$\begin{array}{rcl} \rho(M(L^0), \alpha_{\omega^0}) & = & \rho(M(L), \alpha_\omega) + \sum_{i=1}^m \rho(S^1 \times S^2, \alpha_{z_i}) \\ & = & \rho(M(L), \alpha_\omega). \end{array} \tag{3.2}$$

Let the elements of A be parametrized by the sequence $(a_k)_{k=1}^m$. The link, J, is obtained from L^0 by sliding U_k over the zero framed reverse of L_{a_k} for



Figure 3. Left to right: (1) The two component link L. The map α_ω is trivial on the the meridian of the second component of L. (2) The link L_0 obtained by adding an unknotted component for each ω_k equal to 1. (3) The colored link J is obtained from L_0 by a zero framed handleslide. Proposition 3.3 claims that $\rho(M(L), \alpha_\omega) = \sigma_J(\omega^J)$.

$k = 1, \dots, m$. Kirby calculus (See [13]) reveals that the zero surgeries $M(L^0)$ and $M(J)$ are homeomorphic. The homeomorphism, $\phi : M(L^0) \to M(J)$, between them is given on the level of first homology by

$$\begin{array}{r}H_1(M(L^0)) = \langle \mu_1, \dots \mu_n, \nu_1, \dots, \nu_m \rangle \xrightarrow{\phi_*} \\ H_1(M(J)) = \langle \{\mu_j\}_{j \notin A}, \{\mu^1_{a_k}, \mu^2_{a_k}\}_{a_k \in A} \rangle\end{array}$$

$$\begin{array}{rcl}\mu_j & \mapsto & \mu_j \text{ if } j \notin A \\ \mu_j & \mapsto & \mu^1_j - \mu^2_j \text{ if } j \in A \\ \nu_j & \mapsto & \mu^2_{a_j}\end{array}$$

where μ_j is the meridian of L_j, ν_j is the meridian of U_j and $\mu^1_{a_k}$ and $\mu^2_{a_k}$ are the meridians of the two parallel copies of L_{a_k}. Thus, the following diagram commutes:

$$\begin{tikzcd}H_1(M(L^0)) \arrow[r, "\alpha^0_\omega"] \arrow[d, "\phi_*"'] & U(1) \\ H_1(M(J)) \arrow[ur, "\alpha'"'] & \end{tikzcd}$$

where α' is defined by

$$\begin{array}{rcll}\alpha'(\mu_j) & = & [\omega_j] & \text{if } j \notin A \\ \alpha'(\mu^1_{a_k}) & = & [z_k] & \\ \alpha'(\mu^2_{a_k}) & = & [z_k] & \end{array}$$

Let J^0 denote the $(n+m)$-colored link given by taking all of the components of J to be colored distinctly. By design, $\alpha' : \pi_1(M(J)) \to U(1)$ is realized as $\alpha_{\omega'}$ with $\omega' \in \mathbb{T}^{n+m}_{\mathbb{Q}} \cap \mathbb{T}^{n+m}_{*}$. Thus, equation (3.1) applies and

$$\rho(M(L), \alpha_\omega) = \rho(M(J), \alpha_{\omega'}) = \sigma_{J^0}(\omega').$$

Recall the following property of the Cimasoni-Florens signature:

Proposition (Proposition 2.5, [4]). *Let $L = L(1), \dots, L(n+1)$ be an $(n+1)$-colored link and $L' = L'(1), \dots, L'(n)$ be the n-colored link with $L'(k) = L(k)$ for $k < n$ and $L'(n) = L(n) \cup L(n+1)$. For any $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{T}^n_*$,*

$$\sigma_{L'}(\omega_1, \dots, \omega_n) = \sigma_L(\omega_1, \dots, \omega_n, \omega_n) - \operatorname{lnk}(L_n, L_{n+1})$$

where $\operatorname{lnk}(L_n, L_{n+1})$ is the total linking number between the sublinks L_n and L_{n+1}.

The n-colored link J is obtained from J^0 by identifying the $(a_k)^{\text{th}}$ and $(n+k)^{\text{th}}$ colors. The $(a_k)^{\text{th}}$ and $(n+k)^{\text{th}}$ entries of ω' are both equal to z_k. The n-tuple ω^J is obtained from ω' by deleting the final m terms. The above proposition implies that $\sigma_{J_0}(\omega') = \sigma_J(\omega^J)$ and this completes the proof. $\square$

In light of Proposition 3.3, we make the following extension of $\sigma_L(\omega)$.

Definition 3.4. Let L be a n-component, n-colored link with zero pairwise linking. Let ω be an n-tuple of roots of unity, $A = \{a : \omega_a = 1\}$ and $m = |A|$. Let J be the $(m+n)$-component, n-colored link obtained by replacing $L(a)$ with two zero framed pushoffs of $L(a)$ with opposite orientation from each other and the same color as $L(a)$ for each $a \in A$. Let

$$\omega^J = (\omega^J_1, \dots \omega^J_n) \in \mathbb{T}^n_{\mathbb{Q}} \cap \mathbb{T}^n_*$$

be an n-tuple of roots of unity not equal to 1 with $\omega^J_k = \omega_k$ if $k \notin A$. Then $\widehat{\sigma}_L(\omega) := \sigma_J(\omega^J)$.

Rephrasing Proposition 3.3 in terms of $\widehat{\sigma}_L$, we see that

Theorem 3.5. *For $\omega = (\omega_1, \dots, \omega_n) \in \mathbb{T}^n_{\mathbb{Q}}$, the representation α_ω from $\pi_1(M(L))$ to the subgroup of $U(1)$ given by mapping μ_i to $[\omega_i]$, $\rho(M(L), \alpha_\omega) = \widehat{\sigma}_L(\omega)$.*

We end this section with the proofs of Propositions 3.1 and 3.2.

Proof of Theorem 3.1. Consider the Mayer-Veitoris long exact sequence twisted by α_ω corresponding to the decomposition of W as the union of $E(F)$ together with n disjoint copies of $S^1 \times B^2 \times I$ glued together along n disjoint copies of $S^1 \times S^1 \times I$:

$$\begin{aligned} &\cdots \to \bigoplus_{i=1}^{n} H_2(S^1 \times S^1 \times I; \mathbb{C}_{\alpha_\omega}) \to \\ &H_2(E(F); \mathbb{C}_{\alpha_\omega}) \oplus \left(\bigoplus_{i=1}^{n} H_2(S^1 \times D^2 \times I; \mathbb{C}_{\alpha_\omega})\right) \\ &\to H_2(W; \mathbb{C}_{\alpha_\omega}) \to \bigoplus_{i=1}^{n} H_1(S^1 \times S^1 \times I; \mathbb{C}_{\alpha_\omega}) \to \dots \end{aligned} \tag{3.3}$$

The twisted chain complex for the i^{th} copy of $S^1 \times D^2 \times I \sim S^1$ is chain homotopy equivalent to

$$\mathbb{C} \xrightarrow[1-\omega_i]{} \mathbb{C} \ ,$$

while the twisted chain complex for the i^{th} copy of $S^1 \times S^1 \times I \sim S^1 \times S^1$ is chain homotopy equivalent to

$$\mathbb{C} \xrightarrow[\begin{pmatrix} 0 \\ 1-\omega_i \end{pmatrix}]{} \mathbb{C}^2 \xrightarrow[\begin{pmatrix} 1-\omega_i & 0 \end{pmatrix}]{} \mathbb{C} \ .$$

Both of these chain complexes are acyclic since $\omega_i \neq 1$. Thus, $H_p(S^1 \times S^1 \times I; \mathbb{C}_\omega)$ and $H_p(S^1 \times B^2 \times I; \mathbb{C}_\omega)$ both vanish for all p and the exact sequence (3.3) reveals that $H_2(E(F); \mathbb{C}_\omega) \to H_2(W; \mathbb{C}_\omega)$ is an isomorphism. □

We now prove Proposition 3.2 showing that the unitary ρ-invariants of product 3-manifolds vanish. This will complete the proof of Theorem 3.5.

Proof of Proposition 3.2. Let s denote the element of $H_1(F \times S^1)$ given by the S^1-factor.

Suppose first that $\alpha(s) = [1]$ is the 1×1 identity matrix. In this case α extends over $F \times B^2$, that is the following diagram commutes

$$\begin{tikzcd}
\pi_1(F \times S^1) \arrow[r, "\alpha"] \arrow[d] & U(1) \\
\pi_1(F \times B^2) \arrow[ur, "\alpha"] &
\end{tikzcd}$$

Since $F \times B^2$ deformation retracts to a subset of its boundary it follows that the inclusion induced maps $H_2(F \times S^1) \to H_2(F \times B^2)$ and $H_2(F \times S^1, \mathbb{C}_\alpha) \to H_2(F \times B^2, \mathbb{C}_\alpha)$ are epimorphisms and the twisted and untwisted signatures vanish. Thus, $\rho(F \times S^1, \alpha) = \sigma(F \times B^2, \alpha) - \sigma(F \times B^2) = 0$

W next consider the case that $\alpha(s) \neq [1]$. If F has genus at least two, then let γ be an essential separating simple closed curve. Let W_0 be the 3-manifold given by adding to $F \times [0,1]$ a two handle along $\gamma \times \{1\}$. Then $\partial W_0 = F \sqcup -F_1 \sqcup -F_2$ where F_1 and F_2 have genus strictly less than the genus of F. Since γ is nullhomologous in F, the inclusion induced maps $H_1(F) \overset{\cong}{\to} H_1(W_0)$, and $H_1(F_1) \oplus H_1(F_2) \overset{\cong}{\to} H_1(W_0)$ are isomorphisms. Let $W = W_0 \times S^1$ so that the map $H_1(F \times S^1) \to H_1(W)$ is an isomorphism. Thus, α extends over $\pi_1(W)$ and restricts to representations of $\pi_1(F_1 \times S^1)$ and $\pi_1(F_2 \times S^1)$. Thus,

$$\rho(F \times S^1, \alpha) - \rho(F_1 \times S^1, \alpha) - \rho(F_2 \times S^1, \alpha) = \sigma(W, \alpha) - \sigma(W).$$

Alternately, W_0 can be obtained from $F_1 \sqcup F_2$ by adding a one-handle between different components so that $H_1(W_0, F_1 \sqcup F_2) = H_2(W_0, F_1 \sqcup F_2) =$

0. By the Künneth formula then $H_2(W, F_1 \times S^1 \sqcup F_2 \times S^2) = 0$ so that $H_2(\partial W) \twoheadrightarrow H_2(W)$ is surjective and $\sigma(W) = 0$.

We now compute $\sigma(W, \alpha)$. Consider the Mayer Vietoris long exact sequence

(3.4)

$$\begin{aligned} \dots \to H_2(S^1 \times S^1 \times I; \mathbb{C}_\alpha) \to H_2(F \times S^1; \mathbb{C}_\alpha) \oplus H_2(D^2 \times S^1 \times I; \mathbb{C}_\alpha) \\ \to H_2(W; \mathbb{C}_\alpha) \to H_1(S^1 \times S^1 \times I; \mathbb{C}_\alpha) \to \dots \end{aligned}$$

Since $D^2 \times S^1 \times I \sim S^1$ is a homotopy one complex, it follows that $H_2(D^2 \times S^1 \times I; \mathbb{C}_\alpha) = 0$. The chain complex for $S^1 \times S^1 \times I \sim S^1 \times S^1$ twisted by α is chain homotopic to

$$\begin{tikzcd} \mathbb{C} \arrow[r, "{\begin{pmatrix} 0 \\ 1-\alpha(s) \end{pmatrix}}"'] & \mathbb{C}^2 \arrow[r, "{\begin{pmatrix} 1-\alpha(s) & 0 \end{pmatrix}}"'] & \mathbb{C} \end{tikzcd} .$$

Since $\alpha(s) \neq 1$, this sequence is acyclic and $H_*(S^1 \times S^1 \times I) = 0$. This reduces the sequence (3.4) to the conclusion that $H_2(F \times S^1; \mathbb{C}_\alpha) \to H_2(W; \mathbb{C}_\alpha)$ is an isomorphism so that $\sigma(W, \alpha) = 0$.

Thus, $\rho(F \times S^1, \alpha) - \rho(F_1 \times S^1, \alpha) - \rho(F_2 \times S^1, \alpha) = 0$ where the genera of F_1 and F_2 are each less than the genus of F. An argument inducting on the genus of F will complete the proof once we have proved the Lemma in the case that F is a torus, $\mathbb{T}^2$, or a sphere, S^2.

In the case $F = S^2$, α extends over $S^1 \times B^3$, a homotopy 1-complex. Thus, $\rho(S^2 \times S^1, \alpha) = \sigma(S^1 \times B^3, \alpha) - \sigma(S^1 \times B^3) = 0$.

If $F = \mathbb{T}^2$ then $F \times S^1 = \mathbb{T}^3$ is a 3-torus. Since every finitely generated subgroup of the roots of unity is finite cyclic, α factors as

$$\pi_1(\mathbb{T}^3) \overset{\alpha_0}{\twoheadrightarrow} \mathbb{Z}_p \hookrightarrow U(1).$$

For some basis, $\{b_1, b_2, b_3\}$, for $\pi_1(\mathbb{T}^3) \cong \mathbb{Z}^3$, $\alpha_0(b_1) = \alpha_0(b_2) = 1$ are trivial and $\alpha_0(b_3)$ is a generator of $\mathbb{Z}_p$. Let f_* be the automorphism of $\pi_1(\mathbb{T}^3)$ sending the basis $\{s_1, s_2, s_3\}$ coming from the product structure of $\mathbb{T}^3 = S^1 \times S^1 \times S^1$ to $\{b_1, b_2, b_3\}$. Since $\mathbb{T}^3$ is an Eilenberg-MacClane space, there is a homotopy equivalence $f_0 : \mathbb{T}^3 \to \mathbb{T}^3$ inducing f_*. By [15, Corollary 13.7], f_0 is homotopic to a homeomorphism, f. Thus, we have the following commutative diagram,

$$\begin{tikzcd} \pi_1(\mathbb{T}^3) \arrow[r, "\widehat{\alpha}"] \arrow[d, "f_*"'] & U(1) \\ \pi_1(\mathbb{T}^3) \arrow[ur, "\alpha"'] & \end{tikzcd}$$

where $\widehat{\alpha} = \alpha \circ f_*$. Since ρ-invariants are orientation preserving homeomorphism invariants of 3-manifolds, $\rho(\mathbb{T}^3, \alpha) = \rho(\mathbb{T}^3, \widehat{\alpha})$. Since $\widehat{\alpha}(s_1) = [1]$, $\rho(\mathbb{T}^3, \widehat{\alpha}) = 0$ by the case with which we opened the proof. ☐

We close with an observation which will be necessary in the analysis of the next section.

Corollary 3.6. *For any link L with zero pairwise linking and $\omega \in \mathbb{T}^n_{\mathbb{Q}}$, $\sigma(W, \alpha_\omega) = \widehat{\sigma}_L(\omega)$.*

Proof. By equation (2.1),

$$\sigma(W, \alpha_\omega) = \rho(M(L), \alpha_\omega) - \sum_{i=1}^{n} \rho(\widehat{F}_i \times S^1, \alpha_\omega).$$

By Proposition 3.2, $\sum_{i=1}^{n} \rho(\widehat{F}_i \times S^1, \alpha_\omega) = 0$. Theorem 3.5 now implies the result. ☐

4. Computing L^2-ρ-invariants

We begin by recalling a powerful result of Lück-Schick [24, Theorem 0.1] They prove that if Y is a 4-manifold and $\phi : \pi_1(Y) \to \Lambda$ is a homomorphism to a residually finite group, then $\sigma^{(2)}(Y, \Lambda)$ can be approximated by signatures corresponding to homomorphisms to finite groups. To be precise, if there is a nested sequence of finite index normal subgroups,

$$\Lambda = \Lambda_0 \geq \Lambda_1 \geq \dots,$$

with $\Lambda_0 \cap \Lambda_1 \cap \dots = 0$, then if $p_k : \Lambda \to \Lambda/\Lambda_k$ is the quotient map

$$\sigma^{(2)}(Y, \phi) = \lim_{k \to \infty} \sigma^{(2)}(Y, p_k \circ \phi) \tag{4.1}$$

In the case that Λ is finite, $l^2(\Lambda) = \mathbb{C}[\Lambda]$ and the von Neumann dimension $\dim_{\mathcal{N}\Lambda}$ used to define the L^2-signature gives the same information as the classical dimension theory (See [22, example 1.14]). To be precise, for a finitely generated $\mathbb{C}[\Lambda]$-module V, $\dim_{\mathcal{N}\Lambda}(V) = \frac{1}{|\Lambda|} \dim(V)$, where $\dim(V)$ is the dimension of V as a $\mathbb{C}$-vector space. Unwinding these facts

$$\sigma^{(2)}(Y, \phi) = \frac{1}{|\Lambda|} \sigma(\widetilde{Y}_\phi).$$

where $\widetilde{Y_\phi}$ is the cover of Y corresponding to ϕ

Additionally, in the case that Λ is finite, a classical fact of representation theory says that $\mathbb{C}[\Lambda]$ is the direct sum of irreducible unitary representations of Λ. That is, if $X = \{\alpha : \Lambda \to U(V_\alpha)\}$ is the set of all irreducible unitary representations of Λ, then $\mathbb{C}[\Lambda]$ is isomorphic as a $\mathbb{C}[\Lambda]$-module to $\oplus_{\alpha \in X} V_\alpha^{\dim(V_\alpha)}$. (for example [26, Section 6.2 Proposition 10]). In particular, each V_α is projective and hence flat. Thus,

$$H_2(\widetilde{Y}_\phi) \cong H_2(Y; \mathbb{C}[\Lambda]) \cong H_2(Y; \oplus_{\alpha \in X} V_\alpha^{\dim(V_\alpha)}) \cong \oplus_{\alpha \in X} H_2(Y; V_\alpha)^{\dim(V_\alpha)}.$$

Moreover, the intersection form on $H_2(\widetilde{Y_\phi})$ splits as the direct sum of the intersection forms on $H_2(Y;V_\alpha)$. Thus,

$$\sigma^{(2)}(Y,\phi)=\frac{1}{|\Lambda|}\sigma(\widetilde{Y_\phi})=\frac{1}{|\Lambda|}\sum_{\alpha\in X}\dim(V_\alpha)\sigma(Y,\alpha\circ\phi). \tag{4.2}$$

Between equations (4.1) and (4.2) we see that the unitary signatures of a 4-manifold determine its finite and residually finite L^2-signatures.

Using this mindset together with Corollary 3.6, we express the L^2-signatures of W (from Construction 2.1) as a limit of Riemann sums for an integral of $\widehat{\sigma}_L$. We prove the following theorem:

Proposition 4.1. *Let $A=\langle g_1,\dots,g_n|r_1,\dots,r_m\rangle$ be a finitely generated abelian group. Let W be the 4-manifold of Construction 2.1. Let $\phi: H_1(W)\to A$ be given by sending μ_i to g_i, then*

$$\sigma^{(2)}(W,\phi)=\frac{1}{\lambda(\mathbb{T}_A)}\int_{\mathbb{T}_A}\widehat{\sigma}_L(\omega)d\lambda(\omega) \tag{4.3}$$

where λ is Lebesgue measure on $\mathbb{T}_A$.

Proof. We begin with the case that A is a finite abelian group. In this case equation (4.2) together with the fact that $\mathbb{T}_A$ parametrizes all irreducible representations of A gives that

$$\sigma^{(2)}(W,\phi)=\frac{1}{|A|}\sum_{\omega\in\mathbb{T}_A}\sigma(W,\alpha_\omega).$$

By Corollary 3.6 $\sigma(W,\alpha_\omega)=\widehat{\sigma}_L(\omega)$:

$$\sigma^{(2)}(W,\phi)=\frac{1}{|A|}\sum_{\omega\in\mathbb{T}_A}\widehat{\sigma}_L(\omega). \tag{4.4}$$

When A is finite, $|A|=|\mathbb{T}_A|$, λ is counting measure and integration against λ is summation. After these substitutions, equation (4.4) becomes equation (4.3), completing the proof in the case the A is finite abelian.

Next, for any finitely generated abelian group A, A has a resolution by finite index subgroups $A\geq A_1\geq A_2\geq\dots$ where $A_k=k!A=\langle k!g_1,\dots,k!g_n\rangle$. (Recall that $k!=k(k-1)\dots(3)(2)(1)$ denotes "k-factorial".) The quotient, A/A_k, has a presentation

$$A/A_k=\langle g_1,\dots,g_n,|r_1,\dots,r_m,k!g_1,\dots,k!g_n\rangle$$

and $\mathbb{T}_{A/A_k}=\mathbb{T}_A\cap\{(\omega_1,\dots,\omega_n):\omega_1^{k!}=\dots=\omega_n^{k!}=1\}\subseteq\mathbb{T}^n_{\mathbb{Q}}$.

Thus, by (4.1) and (4.2), for a homomorphism $\phi:\pi_1(W)\to A$ and the projection map $p_k:A\to A/A_k$,

$$\begin{aligned}\sigma^{(2)}(W,\phi) &= \lim_{k\to\infty}\sigma^{(2)}(W,p_k\circ\phi)\\ &= \lim_{k\to\infty}\frac{1}{|\mathbb{T}_{A/A_k}|}\sum_{\omega\in\mathbb{T}_{A/A_k}}\widehat{\sigma}_L(\omega).\end{aligned}$$

This realizes $\sigma^{(2)}(W,A)$ as a limit of Riemann sums for $\frac{1}{\lambda(\mathbb{T}_A)}\int_{\mathbb{T}_A}\widehat{\sigma}_L(\omega)d\lambda(\omega)$. By [4, Theorem 4.1] $\widehat{\sigma}_L$ is piecewise continuous and so Riemann sums converge to the integral. This completes the proof. $\square$

The following theorem shows that a great many $\rho^{(2)}$-invariants of product 3-manifolds vanish. It is much stronger than is needed for our applications. We hope that this result is of independent interest. We delay its proof until subsection 4.1.

Proposition 4.2. *For a closed oriented surface F and any homomorphism to a residually finite group (for example a finitely generated abelian group) $\phi:\pi_1(F\times S^1)\to H$, $\rho^{(2)}(F\times S^1,\phi)=0$.*

As an immediate consequence we see that

Theorem 4.3. *Let $A=\langle g_1\dots g_n|r_1\dots r_m\rangle$ be an abelian group. Let L be an n-component link with zero pairwise linking numbers. Let $\phi:H_1(W)\to A$ be given by sending μ_i to g_i, then*

$$\rho^{(2)}(W,\phi)=\frac{1}{\lambda(\mathbb{T}_A)}\int_{\mathbb{T}_A}\widehat{\sigma}_L(\omega)d\lambda(\omega)$$

where λ is Lebesgue measure on $\mathbb{T}_A$.

Proof. By equation (2.2), $\rho^{(2)}(W,\phi)=\sigma^{(2)}(W,\phi)+\sum\rho^{(2)}(\widehat{F}_i\times S^1,\phi)$. Then by Propositions 4.1 and 4.2,

$$\rho^{(2)}(W,\phi)=\frac{1}{\lambda(\mathbb{T}_A)}\int_{\mathbb{T}_A}\widehat{\sigma}_L(\omega)d\lambda(\omega).$$

$\square$

4.1. The residually finite L^2-ρ-invariants of product manifolds.

We close this section with the proof of Proposition 4.2.

Proof of Proposition 4.2. The von Neumann $\rho^{(2)}$-invariant has the property that for a 3-manifold M, groups A and B, a homomorphism $\phi:\pi_1(M)\to A$ and a monomorphism $\psi:A\hookrightarrow B$, $\rho^{(2)}(M,\phi)=\rho^{(2)}(M,\psi\circ\phi)$. (See [7, Proposition 5.13].) Thus, by replacing H by $\mathrm{im}(\phi)$, we may assume that ϕ is onto.

Let s denote the generator of $\pi_1(S^1)$ so that $\pi_1(F\times S^1)=\pi_1(F)\oplus\langle s\rangle$. Let $\phi(s)=h$. Since s is central in $\pi_1(F\times S^1)$ and ϕ is onto, h is central in H.

The proof proceeds in three steps:

1. We express $\rho^{(2)}(F\times S^1,\phi)$ as a limit of L^2-ρ-invariants corresponding to homomorphisms to finite groups.
2. We reduce the computation of $\rho^{(2)}(F\times S^1,\phi)$ for a homomorphism ϕ from $\pi_1(F\times S^1)$ to a finite group G to the case that $G\cong\mathbb{Z}_n$ is finite cyclic and is generated by $\phi(s)$.
3. Finally we show that $\rho^{(2)}(F\times S^1,\phi)=0$ when the target of ϕ is a finite cyclic group generated by $\phi(s)$.

Step 1: Let Σ be a once puntured torus and $\{a, b\}$ be a symplectic basis for Σ. Let $E = \pi_1(\Sigma)$ be the free group on a and b. Let X be the 4-manifold $F \times \Sigma$. Consider the group,

$$G = \frac{H \oplus E}{h = [a, b]}$$

where $[a, b] = aba^{-1}b^{-1}$. Denote the obvious map $\pi_1(X) = \pi_1(F) \oplus E \to G$ by $\overline{\phi}$. On the boundary, this factors through H. Since h is central in H it does so injectively. That is, the following diagram commutes

$$\begin{tikzcd}
\pi_1(F \times S^1) \arrow[r, "\phi"] \arrow[d] & H \arrow[d, hook] \\
\pi_1(F \times \Sigma) \arrow[r, "\overline{\phi}"] & G
\end{tikzcd}$$

and $\rho^{(2)}(F \times S^1, \phi) = \sigma^{(2)}(F \times \Sigma, \overline{\phi}) - \sigma(F \times \Sigma)$.

We hope to use the behavior of residually finite L^2-signatures to compute $\sigma^{(2)}(F \times \Sigma, \overline{\phi})$. We must show that G is residually finite.

Lemma 4.4. *Let $E = \langle a, b | \rangle$ be the rank two free group with generators a and b. Let H be any residually finite group. Let h be an element of the center of H (i.e. h commutes with every element of H.) Then*

$$G = \frac{H \oplus E}{h = [a, b]}$$

is residually finite.

Proof. We begin by noticing that in G, both a and b commute with $h = [a, b]$, so that $G \cong \frac{H \oplus E'}{h=[a,b]}$ where $E' = \langle a, b | [a, [a, b]] = [b, [a, b]] = 1\rangle$ is isomorphic to the 3-dimensional integral Heisenberg group. That is, the multiplicative group of 3×3 upper triangular integral matrices with ones on the main diagonal,

$$E' \cong \left\{ \begin{pmatrix} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix} : \alpha, \beta, \gamma \in \mathbb{Z} \right\}. \tag{4.5}$$

Let $e(\alpha, \beta, \gamma) = \begin{pmatrix} 1 & \alpha & \gamma \\ 0 & 1 & \beta \\ 0 & 0 & 1 \end{pmatrix}$. The isomorphism is given by sending $a \mapsto e(1, 0, 0)$, $b \mapsto e(0, 1, 0)$. As a consequence, $[a, b] \mapsto e(0, 0, 1)$. For any integer n, let $\phi_n : E' \to E_n \leq Gl(3, \mathbb{Z}_n)$ be the homomorphism given by reduction of entries mod n. In this finite quotient, $\phi_n([a, b])$ is order n.

Let g be a nontrivial element of G. We will find a finite quotient of G in which g is still non-trivial. If we can do so we will conclude that G is residually finite.

Notice that g is represented by an equivalence class $(x \oplus y)$, $x \in H$, $y \in E'$. If y does not sit in the cyclic subgroup generated by $[a,b]$ in E' (which is normal), then $(x \oplus y)$ is nonzero in the quotient, $q : G \to \frac{E'}{([a,b])} \cong \mathbb{Z}^2$, whose target is residually finite. In some finite quotient of $\mathbb{Z}^2$, $q(g) = q(x \oplus y)$ is nonzero.

Alternately, if $y = [a,b]^m$ for some $m \in \mathbb{Z}$, then $g = (x \oplus [a,b]^m) = (xh^m \oplus 1)$ in G. If $(x \oplus y)$ is nontrivial, then $x' = xh^m$ is nontrivial in H, which by assumption is residually finite. Let $f : H \to \overline{H}$ be a homomorphism to a finite group with $f(x') \neq 1$. Let n be the order of $f(h)$ in this group. The direct sum $f \oplus \phi_n : H \oplus E' \to \overline{H} \oplus E_n$ passes to a homomorphism $F : G \to \frac{\overline{H} \oplus E_n}{f(h) = \phi_n([a,b])}$.

Consider the resulting commutative diagram

$$\begin{tikzcd}
H \arrow[r] \arrow[d, "f"] & H \oplus E' \arrow[r] \arrow[d, "f \oplus \phi_n"] & \dfrac{H \oplus E'}{h = [a,b]} \arrow[d, "F"] \\
\overline{H} \arrow[r] & \overline{H} \oplus E_n \arrow[r] & \dfrac{\overline{H} \oplus E_n}{f(h) = \phi_n([a,b])}
\end{tikzcd}$$

Since h is central in H, $f(h)$ is central in $\overline{H}$ and the composition along the bottom row is injective. Since by assumption, $f(x') \neq 1$, We conclude that $F(g) = F(x' \oplus 1) \neq 1$.

Thus, any nontrivial element of G is nontrivial in some finite quotient and G is residually finite. This completes the proof.

□

So, G is residually finite. Let $G \geq G_1 \geq G_2 \geq \dots$ be a finite resolution for G and $p_k : G \to G/G_k$ be the quotient map.

$$\begin{aligned}
\rho^{(2)}(F \times S^1, \phi) &= \sigma^{(2)}(F \times \Sigma, \overline{\phi}) - \sigma(F \times \Sigma) \\
&= \lim_{k \to \infty} \left(\sigma^{(2)}(F \times \Sigma, p_k \circ \overline{\phi}) - \sigma(F \times \Sigma) \right) \\
&= \lim_{k \to \infty} \rho(F \times S^1, p_k \circ \overline{\phi})
\end{aligned}$$

Since $p_k \circ \overline{\phi} : \pi_1(F \times S^1) \to G/G_k$ is a homomorphism to a finite group for each k, it remains only to prove the theorem in the case that H is finite.

Step 2: Assuming that H is a finite group, h^n is trivial in H for some n. Let P be the n times punctured sphere. The group, $\pi_1(P)$ is isomorphic to $\langle s_1 \dots s_n | s_1 s_2 \dots s_n = 1 \rangle$. Consider the homomorphism

$$\begin{aligned}
&\overline{\phi} : \pi_1(F \times P) \cong \pi_1(F) \oplus \langle s_1 \dots s_n | s_1 s_2 \dots s_n = 1 \rangle \to H \\
&\overline{\phi}(f) = \phi(f) \text{ if } f \in \pi_1(F) \\
&\overline{\phi}(s_i) = h \text{ for } i = 1, \dots, n
\end{aligned}$$

Notice that $\partial(F \times P) = F \times \partial P$ is given by n copies of $F \times S^1$. On each of these boundary components, $\overline{\phi}$ restricts to ϕ. Thus, $n\rho(F \times S^1, \phi) = \sigma^{(2)}(F \times P, \overline{\phi}) - \sigma(F \times P)$. By a Künneth theorem based argument, the map $H_2(F \times \partial P) \to H_2(F \times P)$ is surjective and

$$\sigma(F \times P) = 0. \tag{4.6}$$

Since H is finite, the L^2-signature and the signature of the $\overline{\phi}$-cover agree,

$$\sigma^{(2)}(F \times P, \overline{\phi}) = \frac{1}{|H|}\sigma(\widetilde{F \times P}_{\overline{\phi}}).$$

In order to compute $\sigma(\widetilde{F \times P}_{\overline{\phi}})$, we first study the cover of $F \times P$ corresponding to the composition $\psi : \pi_1(F \times P) \xrightarrow{\overline{\phi}} H \to \frac{H}{\langle h \rangle}$. Again, h is central in H, so the cyclic subgroup it generates is normal in H and this quotient makes sense. Since ψ is trivial on $\pi_1(P)$, the corresponding cover is given by the product of a (compact) cover of F with P, $\widetilde{F}_\psi \times P$. A compact cover of a closed oriented surface is still a closed oriented surface.

Since ψ factors through ϕ, the covering map corresponding to ϕ factors through the covering map corresponding to ψ. Consider the resulting tower of covers.

$$\begin{tikzcd}
\widetilde{F \times P}_{\overline{\phi}} \arrow[dd] \arrow[dr, "\alpha"] & \\
& \widetilde{F}_\psi \times P \arrow[dl] \\
F \times P &
\end{tikzcd}$$

The group of deck translations of α is isomorphic to

$$\frac{\pi_1(\widetilde{F}_\psi \times P)}{\alpha_*[\widetilde{F \times P}_{\overline{\phi}}]} \cong \frac{\ker(\psi)}{\ker(\overline{\phi})} \cong \overline{\phi}[\ker(\psi)] = \ker\left(H \to \frac{H}{\langle h \rangle}\right) = \langle h \rangle \cong \mathbb{Z}_n$$

Thus, $\widetilde{F \times P}_{\overline{\phi}}$ is a finite cyclic cover of $\widetilde{F}_\psi \times P$. It remains then only to show that for a closed oriented surface $F' = \widetilde{F_\psi}$ and a homomorphism ϕ' from $\pi_1(F' \times P) = \pi_1(F') \oplus \langle s_1 \dots s_n | s_1 s_2 \dots s_n = 1\rangle$ to a finite cyclic group $\mathbb{Z}_n$ which sends each s_i to the same generator of $\mathbb{Z}_n$, the signature of the resulting cover is zero.

Consider the following chain of equalities:

$$\begin{aligned}
\sigma(\widetilde{F' \times P}_{\phi'}) &= n\sigma^{(2)}(F' \times P, \phi') \\
&= n(\sigma^{(2)}(F' \times P, \phi') - \sigma(F' \times P)) \\
&= n\rho^{(2)}(F' \times S^1, \phi').
\end{aligned}$$

The first follows from equation (4.2), the second from (4.6). The third is by the definition of $\rho^{(2)}$-invariants. It remains only to show that for a homomorphism $\phi' : \pi_1(F' \times S^1) = \pi_1(F') \oplus \langle s \rangle \to \mathbb{Z}_n$ sending s to a generator of $\mathbb{Z}_n$, $\rho^{(2)}(F' \times S^1, \phi') = 0$.

Step 3: The group $\pi_1(F')$ has presentation

$$\pi_1(F') \cong \langle a_1, b_1, \ldots, a_g, b_g | [a_1, b_1] \ldots [a_g, b_g] \rangle$$

where g is the genus of F'. Since $\phi'(s)$ generates $\mathbb{Z}_m$, $\phi(a_i) = \phi'(s)^{p_i}$ for some $p_i \in \mathbb{Z}$. Consider the automorphism of $\pi_1(F' \times S^1)$ given by $s \mapsto s$, $a_i \mapsto a_i s^{-p_i}$, $b_i \mapsto b_i$. Similarly to the analysis in the proof of Proposition 3.2, this map is induced by a homeomorphism $\Phi : F' \times S^1 \to F' \times S^1$.

Thus, $\rho^{(2)}(F' \times S^1, \phi') = \rho^{(2)}(F' \times S^1, \phi' \circ \Phi_*)$, and $\phi_0 := \phi' \circ \Phi_*$ is trivial on every a_i. Let V be a handle body bounded by F' given by adding 2-handles along every a_i. Since ϕ_0 is trivial on a_i, ϕ_0 extends to a homomorphism $\overline{\phi_0} : \pi_1(V \times S^1) \to \mathbb{Z}_n$. There is a deformation retraction from V to a subset of F'. This implies that $H_2(F' \times S^1) \twoheadrightarrow H_2(V \times S^1)$ and $H_2(F' \times S^1; \mathbb{C}[\mathbb{Z}_n]) \twoheadrightarrow H_2(V \times S^1; \mathbb{C}[\mathbb{Z}_n])$ are both epimorphisms. Thus, both the classical intersection form and the L^2-intersection form of $V \times S^1$ are zero and the signature and L^2-signature both vanish. Thus,

$$\rho^{(2)}(F' \times S^1, \phi_0) = \sigma^{(2)}(V \times S^1, \overline{\phi_0}) - \sigma(V \times S^1) = 0,$$

completing the proof.

□

5. Application: The linear independence of the twist knots which are algebraically of order 2

In this section, we study those twist knots (depicted in Figure 1) which are algebraically of order 2 using the tools established in this paper.

We recall some concepts from algebraic concordance. Let K be an algebraically slice knot with a genus g Seifert surface F. Then, by the definition of algebraic concordance, there exists a nonseparating collection of g simple closed curves on F, $L = L_1 \ldots L_g$, for which the Seifert form of F vanishes. That is, $\operatorname{lnk}(L_i, L_j^+) = 0$ for all i, j, where L_j^+ is the result of pushing L_j off of F in the positive normal direction and lnk is the the linking number. Using language from [5], L is called a **derivative** of K.

For a simple closed curve, γ, on F, a curve m in the complement of F is called a **meridian for the band on which γ sits** if m bounds a disk which intersects F in a single arc and intersects γ in a single point. All of these intersections must be transverse.

In this section we consider ρ^0, the ρ-invariant of $M(L)$ corresponding to abelianization. To be precise, if L is an n-component link with zero pairwise linking and $\phi : \pi_1(M(L)) \to \mathbb{Z}^n$ is the abelianization homomorphism then $\rho^0(L) := \rho^{(2)}(M(L), \phi)$. By Theorem 4.3 This is given by the integral of the Cimasoni-Florens signature of L over all of $\mathbb{T}^n$.

In [10] the author considers a particular metabelian ρ-invararant of knots, ρ^1 and proves the following theorems. Stronger theorems are proven in that paper, but these are as strong as required in this paper.

Theorem ([10], Corollary 4.4)**.** *If $K_1, \dots K_n$ are knots with distinct prime Alexander polynomials, vanishing ρ^0-invariants and nonzero ρ^1-invariants, then $\{K_i\}$ is linearly independent in the knot concordance group.*

Theorem ([10], Theorem 5.6)**.** *Suppose that K is a genus 1 knot and that $K\#K$ is algebraically slice. Let L be a two component derivative of $K\#K$. If the components of L together with the meridians of the bands on which L sit form a $\mathbb{Z}$-linearly independent set in the Alexander module of $K\#K$, then*

$$|2\rho^1(K) - \rho^0(L)| \leq 1.$$

As a first step in applying the tools of this paper to those twist knots which are algebraically of order 2 we find for every n such that $T_n\#T_n$ is algebraically slice a derivative of $T_n\#T_n$. Namely we find that the link of Figure 4 is a derivative.



FIGURE 4. For $n = a^2 - a + b^2$, the link $L_{a,b}$ is a derivative for T_n The curves m_1 and m_2 are meridians about the bands on which the components of L sit.

Lemma 5.1. *The n-twist knot T_n is algebraically of order exactly 2 if and only if there are positive integers a and b such that $n = a^2 - a + b^2$ but there does not exist an integer c with $n = c^2 - c$. For such an n, the link $L_{a,b}$, depicted in Figure 4, is a derivative for $T_n\#T_n$. Moreover the meridians, m_1 and m_2, of the bands on which the components of $L_{a,b}$ sit are $\mathbb{Z}$-linearly independent of the components of L in the Alexander module of $T_n\#T_n$.*

Proof. According to Levine [18, Corollary 23], $T_n\#T_n$ is algebracially slice if and only if $4n+1$ has no odd multiplicity prime factors congruent to 3 mod 4. Under these conditions an elementary fact from number theory (see [3, Theorem 12.3], for example) provides integers a_0 and b_0 with $a_0^2+b_0^2 = 4n+1$. Since $4n+1$ is odd it must be that $a_0 = 2a-1$ is odd while $b_0 = 2b$ is even. Under these conditions this equation reduces to $a^2 - a + b^2 = n$ as was claimed.



FIGURE 5. $\gamma_1, \dots, \gamma_4$ form a basis for the first homology of a Seifert surface for $T_n \# T_n$. The meridians for γ_i, $\mu_1, \dots, \mu_4$ form a generating set for the Alexander module of $T_n \# T_n$.

We now prove that the link $L_{a,b}$ is a derivative. The first homology of F is a free abelian group with basis given by the curves $\gamma_1, \gamma_2, \gamma_3, \gamma_4$ in Figure 5. With respect to this basis the components of $L_{a,b}$ represent the classes in $H_1(F)$ represented by the vectors

$$v_1 = \left[\begin{array}{cccc} 1 & a & 0 & b \end{array} \right]^T, v_2 = \left[\begin{array}{cccc} 0 & b & 1 & (1-a) \end{array} \right]^T$$

while the Seifert form of F has the matrix

$$V = \left[\begin{array}{cccc} n & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & n & 1 \\ 0 & 0 & 0 & -1 \end{array} \right]$$

A computation (remembering that $n = a^2 - a + b^2$) verifies that $v_i^T V v_j = 0$ for $i, j \in \{1, 2\}$. Thus, $L_{a,b}$ is a derivative.

In order to address the linear independence claim we first recall some classical knot theory facts. A good reference is [25, Chapter 8, Section C]. The Alexander module, $A_0(T_n \# T_n)$, is the $\mathbb{Q}[t, t^{-1}]$-module generated by $\mu_1, \mu_2, \mu_3, \mu_4$ with presentation matrix $V - tV^T$. With respect to this presentation, $m_1 = \mu_1$ and $m_2 = \mu_3$ correspond to the first and third generators and γ_i corresponds to the i^{th} column of V.

Performing the presentation calculus needed to unwind this yeilds

$$A_0(T_n \# T_n) \cong \left(\frac{\mathbb{Q}[t, t^{-1}]}{(nt^2 - (2n+1)t + n)} \right) \oplus \left(\frac{\mathbb{Q}[t, t^{-1}]}{(nt^2 - (2n+1)t + n)} \right)$$

with generators m_1 and m_2 and that the components of $L_{a,b}$ (represented by v_1 and v_2) correspond to

$$\begin{array}{rcl} v_1 & \mapsto & (n + a(1 - n + nt))m_1 + b(1 - n + nt)m_2 \\ v_2 & \mapsto & b(1 - n + nt)m_1 + (1 + (1 - a)(1 - n + nt))m_2. \end{array}$$

It is straightforward to see that images of m_1, m_2, v_1 and v_2 form a $\mathbb{Q}$-linearly independent set. This completes the proof. □

Thus, in light of [10, Theorem 5.6] and Theorem 4.3, in order to show that $\rho^1(T_n) \neq 0$, it suffices to show that the integral of the Cimasoni-Florens signature of the link $L_{a,b}$ over $\mathbb{T}^2$ is greater than 1 in absolute value.

5.1. **Tools for computing the Cimasoni-Florens signature.** In [4], Cimason and Florens demonstrate the effect that crossing change and smoothing (see Figure 6) have on the signature function. We make use of these results in order to ease the approximation of the ρ^0-invariant. Observe that for a link with non-zero linking numbers or a colored link with multiple components of a single color, we have demonstrated no relationship between the integral of the Cimasoni-Florens signature function. Since we will need to discuss the integral of the Cimasoni-Florens signature of such links if we are to make use of the moves in Figure 6 we make up some notation. For an arbitrary n-colored link L, the R**-invariant** of L, $R(L)$ is defined as the integral of $\widehat{\sigma}$ over the n-dimensional unit torus with respect to normalized Lebesgue measure:

$$R(L) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} \widehat{\sigma_L}(\omega) d\lambda(\omega)$$



Figure 6. Moves which change the Cimasoni-Florens signature by at most one. The move on the left is understood to preserve orientations and be between arcs of the same color. The move of the right is understood to be between arcs of different colors

Cimasoni and Florens in [4, Proposition 5.1] show that if colored links L and L' differ by either of the the moves indicated in Figure 6 then for all ω, $\sigma_L(\omega)$ and $\sigma_{L'}(\omega)$ differ by 0 or 1 or -1 depending on the Conway potential function of L and L'. Integrating this result gives that $R(L)$ and $R(L')$ differ by at most 1.

The addition of a band between two arcs of the same color (as in Figure 7) can be realized by a single smoothing. Thus, if L and L' differ by either a band addition or a crossing change between arcs of the same color, then $R(L)$ and $R(L')$ differ by at most 1. Additionally, [4, Proposition 2.12] shows that if this band runs between split sublinks of L, then $\sigma_L = \sigma_{L'}$. Summarizing these results,



FIGURE 7. A band addition can be realized via a smoothing

1. If colored links L and L' differ by a crossing change between arcs of different colors, then $|R(L) - R(L')| \leq 1$
2. If colored links L and L' differ by a smoothing a crossing between arcs of the same color, then $|R(L) - R(L')| \leq 1$
3. If colored links L and L' differ by the addition of a band between arcs of the same color, then $|R(L) - R(L')| \leq 1$. If this band joins split sublinks of L then $R(L) = R(L')$.

The local move on colored links depicted in Figure 8 will be relevant in our analysis of $L_{a,b}$. We provide bounds on how it changes the R-invariant.



FIGURE 8. The move considered in Proposition 5.2 realized via band summing with the link V. The two different bands of arcs are assumed to be of different colors.

Proposition 5.2. *If the colored links* L *and* L' *differ by the move depicted in Figure 8 with* a *strands of one color and* b *strands of a different color then* $|R(L) - R(L')| \leq a + b - 1$

Proof. As is shown in Figure 8, L' is given by taking the split union of L with the two color link V depicted in Figure 9 and adding $a{+}b$ bands, with the first band being between split sublinks. Since the signature invariant adds under split union, by [4, Proposition 2.12], it follows that $|R(L') - R(L) - R(V)| \leq a + b - 1$, where V is the link depicted in Figure 9.

In order to compute $R(V)$, we notice that the result of reversing the orientation all of the components of V of one color is isotopic the mirror image of V, so that by [4, Proposition 2.10 and Corollary 2.11], $\sigma_V(\omega_1, \omega_2^{-1}) = -\sigma_V(\omega_1, \omega_2)$. Since $\omega \mapsto \omega^{-1}$ is a measure preserving transformation of $\mathbb{T}^1$, this implies that $R(V) = 0$ and completes the proof. $\square$



FIGURE 9. The link V and an isotopy of it

5.2. **Performing the computation for $L_{a,b}$.** In this section we perform the promised computation of $\rho^0(L_{a,b}) = R(L_{a,b})$. We begin by isotoping $L_{a,b}$ to the link in Figure 10.



FIGURE 10. The link $L_{a,b}$ after an isotopy.

By adding $2b$ bands to this link, b of each color, the link $L_{a,b}$ is reduced to the split union of two links, $L^1_{a,b}$ depicted in Figure 11. Thus, $|\rho^0(L_{a,b}) - R(L^1_{a,b})| \leq 2b - 1$.



FIGURE 11. The link $L^1_{a,b}$ and an isotopy of it.

By changing b crossings between components of different colors $L^1_{a,b}$ becomes the link $L^2_{a,b}$ of Figure 12 so that $|R(L^1_{a,b}) - R(L^2_{a,b})| \leq b$ and $|\rho^0(L_{a,b}) - R(L^2_{a,b})| \leq 3b - 1$

The link $L^2_{a,b}$ is the result of performing two band twist moves (as in Figure 8) starting with the link $L^3_{a,b}$ (which now consists of 4 split factors



FIGURE 12. The link $L^2_{a,b}$ and some isotopies of it. The B denotes the move in Figure 8.

each of which is a one color torus link) of Figure 13. The leftmost twist involves a strands of one color and b of the other. The rightmost involves $a-1$ of one color and b of the other. Thus, $|R(L^2_{a,b}) - R(L^3_{a,b})| \leq (a+b-1) + (a-1+b-1)$, and $|\rho^0(L_{a,b}) - R(L^3_{a,b})| \leq 5b + 2a - 4$.

Finally, since by [4, Proposition 2.13] Cimasoni-Florens signature adds under split union,

$$R(L^3_{a,b}) = R(T(a, 1-a)) + R(T(1-a, a)) + 2R(T(b, -b)),$$

where $T(a, b)$ is the one color (a, b)-torus link. In the case of knots, the Cimasoni-Florens signature agrees with the Tristram-Levine signature. The integral of the Tristram-Levine signature of torus knots is computed by Borodzic [2] and independently by Collins [9]. In [2], Borodzic also computes the integral of the one colored signature of the (b, b)-torus link, which is the mirror image of the $(b, -b)$-torus link. Specifying their results to our setting,

$$\begin{aligned} R(T(a, 1-a)) &= R(T(1-a, a)) = \frac{(a+1)(a-2)}{3} \\ R(T(b, -b)) &= \frac{(b-1)^2}{3} \end{aligned}$$



FIGURE 13. The link $L^3_{a,b}$ consists of the split union of two one color $(b, -b)$-torus links, a $(a-b, b-a+1)$-torus knot and a $(a+b-3, -a-b+2)$-torus knot.

Putting all of this together,

$$\begin{aligned} \rho^0(L_{a,b}) &\geq \frac{2(a+1)(a-2)+2(b-1)^2}{3} - (2a+5b-4) \\ &= \frac{2a^2+2b^2-8a-19b+10}{3} \end{aligned} \tag{5.1}$$

Since we are interested in when $\rho^0(L_{a,b}) > 1$, we ask when

$$f(a,b) := 2a^2+2b^2-8a-19b+7 > 0$$

By observing that $f(a,b)$ grows quadratically in both a and b, we immediately see that

Proposition. *For all but finitely many a and b, $f(a,b) > 0$ so that $\rho^0(L_{a,b}) > 1$ and for $n = a^2 - a + b^2$, $\rho^1(T_n) > 0$. Thus, a set containing all but finitely many twist knots of algebraic order 2 is linearly independent in $\mathcal{C}$.*

Next, We use a computer to determine $\{n : n = a^2 - a + b, a > 0, b > 0, f(a,b) > 0\}$. Combining this result with the computation in [10, Corollary 6.2] showing that $\rho^1(T_n) > 0$ for n of the form $n = x^2 + x + 1$ with $x > 1$ gives the following result. (The knot called T_n in this paper is the mirror image of what is called T_{-n} in [10])

Theorem 5.3. *The set containing all of the twist knots T_n which are algebraically of order 2 is linearly independent with the following 39 possible exceptions:*

$$\begin{aligned} n \;=\; & 1,3,4,9,10,11,15,16,18,22,24,25,27,28,29,34,36,37,38, \\ & 39,45,48,49,51,55,58,61,64,66,67,69,70,78,79,83,84,87, \\ & 93,101. \end{aligned}$$

One can do better by noticing the following consequence of the results of [10].

Proposition 5.4. *Let A be a set of knots with distinct prime Alexander polynomials, vanishing ρ^0-invariants and nonvanishing ρ^1-invariants. Let B be a linearly independent set of knots with vanishing ρ^0-invariants and prime Alexander polynomials distinct from the Alexander polynomials of the elements of A, then $A \cup B$ is linearly independent.*

Proof. The proof relies on a family von Neumann ρ-invariants indexed by the polynomial $p(t)$ defined in [10], $\rho^1_{p(t)}$. The following proposition reduces them in many cases to ρ^0 or ρ^1.

Proposition ([10] Proposition 3.4)**.** *Let $\Delta(t)$ be the Alexander polynomial of a knot K. Then*

1. $\rho^1_{\Delta(t)}(K) = \rho^1(K)$.
2. *If* $(p, \Delta) = 1$, *then* $\rho^1_{p(t)}(K) = \rho^0(K)$.

Moreover, by [10, Theorem 4.1 and Proposition 3.5], ρ^1_p is a homomorphism when restricted to the subgroup of $\mathcal{C}$ generated by knots with prime Alexander polynomials.

Suppose that some linear combination $\left(\underset{K\in A}{\#} l_K K\right) \# \left(\underset{J\in B}{\#} m_J J\right)$ is slice. Let p be the Alexander polynomial of some $K_0 \in A$. Since ρ^1_p is a homomophism, it follows that

$$0 = \sum_{K\in A} l_K \rho^1_p(K) + \sum_{J\in B} m_J \rho^1_p(J) \tag{5.2}$$

Next, by [10, Proposition 3.4], and since ρ^0 vanishes for all of the knots in A and B, equation (5.2) reduces to $0 = l_{K_0}\rho^1(K_0)$. Since $\rho^1(K_0) \neq 0$, it follows that $l_{K_0} = 0$.

Since K_0 was arbitrarily chosen in A, it follows that every l_K vanishes. Finally, since B is assumed to be linearly independent in $\mathcal{C}$, this implies that every $m_J = 0$ and we conclude that $A \cup B$ is linearly independent. □

In [27, Corollary 1.2], Tamulis finds that $\{T_n : 4n+1 \text{ is prime}\}$ is linearly independent in $\mathcal{C}$. Hence, we can remove all elements which satisfy this condition from the set of possible counter-examples of Proposition 5.3.

Theorem 5.5. *The set containing all of the twist knots T_n which are algebraically of order 2 is linearly independent with the following 12 possible exceptions:*

$$n = 1, 11, 16, 29, 36, 38, 51, 55, 61, 66, 83, 101.$$

References

[1] M. F. Atiyah, V. K. Patodi, and I. M. Singer. Spectral asymmetry and Riemannian geometry. II. *Math. Proc. Cambridge Philos. Soc.*, 78(3):405–432, 1975.

[2] Macief Borodzik. A ρ-invariant of iterated torus knots. *Proceedings of Symposia in Pure Mathematics*, 82, 2011.

[3] David M. Burton. *Elementary number theory.* Allyn and Bacon Inc., Boston, Mass., 1980. Revised printing.

[4] David Cimasoni and Vincent Florens. Generalized Seifert surfaces and signatures of colored links. *Transactions of the American Math Society*, 360(3), March 2008.

[5] Tim D. Cochran, Shelly Harvey, and Constance Leidy. Derivatives of knots and second-order signatures. *Algebr. Geom. Topol.*, 10(2):739–787, 2010.

[6] Tim D. Cochran, Shelly Harvey, and Constance Leidy. Primary decomposition and the fractal nature of knot concordance. *Math Annalen*, 2010. DOI: 10.1007/s00208-010-0604-5.

[7] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Knot concordance, Whitney towers and L^2-signatures. *Ann. of Math. (2)*, 157(2):433–519, 2003.

[8] Tim D. Cochran, Kent E. Orr, and Peter Teichner. Structure in the classical knot concordance group. *Comment. Math. Helv.*, 79(1):105–123, 2004.

[9] Julia Collins. The $L^{(2)}$-signature of torus knots. preprint avilable at http://front.math.ucdavis.edu/1001.1329.

[10] Christopher William Davis. Von Neumann rho invariants as obstructions to torsion in the topological knot concordance group. *Algebraic and Geometric Topology*, pages 753–789, 2012. DOI: 10.2140/agt.2012.12.753.

[11] Stefan Friedl. Eta invariants as sliceness obstructions and their relation to Casson-Gordon invariants. *Algebr. Geom. Topol.*, 4:893–934, 2004.

[12] Stefan Friedl. L^2-eta-invariants and their approximation by unitary eta-invariants. *Math. Proc. Cambridge Philos. Soc.*, 138(2):327–338, 2005.

[13] Robert E. Gompf and András I. Stipsicz. 4-*manifolds and Kirby calculus*, volume 20 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1999.

[14] Shelly L. Harvey. Homology cobordism invariants and the Cochran-Orr-Teichner filtration of the link concordance group. *Geom. Topol.*, 12(1):387–430, 2008.

[15] John Hempel. *3-Manifolds*. AMS Chelsea Publishing, Providence, RI, 2004. Reprint of the 1976 original.

[16] Se-Goo Kim. Polynomial splittings of Casson-Gordon invariants. *Math. Proc. Cambridge Philos. Soc.*, 138(1):59–78, 2005.

[17] Se-Goo Kim and Taehee Kim. Polynomial splittings of metabelian von Neumann rho-invariants of knots. *Proc. Amer. Math. Soc.*, 136(11):4079–4087, 2008.

[18] J. Levine. Invariants of knot cobordism. *Invent. Math. 8 (1969), 98–110; addendum, ibid.*, 8:355, 1969.

[19] Paolo Lisca. Sums of lens spaces bounding rational balls. *Algebr. Geom. Topol.*, 7:2141–2164, 2007.

[20] R. A. Litherland. Cobordism of satellite knots. In *Four-manifold theory (Durham, N.H., 1982)*, volume 35 of *Contemp. Math.*, pages 327–362. Amer. Math. Soc., Providence, RI, 1984.

[21] Charles Livingston and Swatee Naik. Obstructing four-torsion in the classical knot concordance group. *J. Differential Geom.*, 51(1):1–12, 1999.

[22] Wolfgang Lück. L^2-invariants from the algebraic point of view. In *Geometric and cohomological methods in group theory*, volume 358 of *London Math. Soc. Lecture Note Ser.*, pages 63–161. Cambridge Univ. Press, Cambridge, 2009.

[23] Wolfgang Lück and Thomas Schick. Various L^2 signatures and a topological L^2 signature theorem. *High-dimensional manifold topology*, pages 362–399, 2003.

[24] Wolfgang Lučk and Thomas Schick. Approximating l^2-signatures by their compact analogues. *Forum Mathematicum*, 17:31–65, 2005.

[25] Dale Rolfsen. *Knots and links*, volume 7 of *Mathematics Lecture Series*. Publish or Perish Inc., Houston, TX, 1990. Corrected reprint of the 1976 original.

[26] Jean-Pierre Serre. *Linear representations of finite groups*. Springer-Verlag, New York, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.

[27] Andrius Tamulis. Knots of ten or fewer crossings of algebraic order 2. *J. Knot Theory Ramifications*, 11(2):211–222, 2002.

Department of Mathematics, Rice UNIVERSITY
E-mail address: `youremailaddress@math.uh.edu`